



Nº PATENTE	SOLICITANTE	PAÍS	TÍTULO
WO2010082832	NOFIMA MARIN	Noruega	Improve texture quality fish fillet select salmon add individual mixture purification free bioactive amino acid diet.
WO2010068990	SHOALHAVEN STARCHES PTY LTD	Australia	Protein-rich material useful in aquaculture feed for providing complementary nutrition to animals such as halibut, eel, crustaceans and mollusks, prepared as composite material using wheat protein ingredient or lupin protein ingredient.
TW201000632	YANG SHERNG BIOTECHNOLOGY CO LTD	Taiwan	Peptide for promoting gonad maturity, mating, and spawning of crustaceans - which promotes gonad maturity, mating and spawning of the crustaceans.
KR20100048509	DONGBU HITEK CO LTD	Corea	Liquid composition for the exterior parasite relief of fishes and preventing gill monogenean disease, comprises praziquantel, tocopheryl derivative as antioxidant and organic solvent.
KR100937037B	DOOSAN ECOBIZNET CO LTD	Corea	Formulation for treating microbial infection in fish, contains vitamin B1, sodium dodecylsulfate, kaolin, sodium lignin sulfonate, diaryl dimethyl ammonium, organopolysiloxane, butylene glycol and glycerin fatty acid ester.
KR100952344B	CHOI G KIM G KIM J	Corea	Preparing flatfish feed useful for enhancing immunity, and preventing and treating microbial infection and dyspepsia, by heating livefood with seed culture, fermenting heated livefood mixture and preparing fermented feed into pellets.
CN101773209	ZHEJIANG KESHENG FEED CO LTD	China	Feed useful for promoting growth of shellfish, comprises protein feed, energy feed, mineral, vitamin and Bacillus preparation.
CN101756063	HONG F	China	Feed additive useful for improving growth environment of breeding fish and shrimp, comprises zeolite, shell powder, calcium hydrophosphate, rhizoma Smilacis glabrae, honeysuckle and Radix stemonae.
CN101731167	CHINA SEAFOOD TECHNOLOGY ACAD HUANGHAI S	China	Individual marking method for selective breeding of turbot, involves putting fish in seawater pool and in medical bathing with florfenicol for specific hours for three days, where concentration of florfenicol is specific parts per million.